

COMPOSITIONAL, PHYSICO-MECHANICAL AND FUNCTIONAL
PROPERTIES OF TWO GHANAIAN COWPEA
(*Vigna unguiculata*) VARIETIES

Tawiah ER^{1,3}, Akonor PT^{2,3*}, Johnson P-N T³, Oduro-Yeboah C^{2,3},
Idun-Acquah NN², Mingle C¹ and TK Tengey



ER Tawiah

*Corresponding author email: papatoah@gmail.com

¹Food and Drugs Authority, Box CT 2783, Accra, Ghana

² Council for Scientific and Industrial Research- Food Research Institute, Box M.20, Accra, Ghana

³ Department of Agro-processing Technology and Food Biosciences, CSIR-College of Science and Technology, Box M.20, Accra, Ghana

⁴ Council for Scientific and Industrial Research- Savanna Agricultural Research Institute, Box 52, Nyankpala, Ghana



ABSTRACT

Cowpea (*Vigna unguiculata*) is an important legume cultivated in many parts of Africa and contributes a significant amount of plant-based protein to human diets. There are many varieties of cowpeas, and these have varying seed characteristics such as shape, size, colour and nutritional properties. The compositional, physico-mechanical and functional properties of two varieties of cowpea seeds were studied in order to establish their full food-use potentials as well as key parameters for their postharvest machinery for processing, handling, and storage. The proximate composition of the two cowpea cultivars was determined using standard methods. Physico-mechanical analyses carried out were the hundred-seed weight and volume, bulk density, true density, size, aspect ratio, surface area, equivalent diameter, porosity and angle of repose. Functional properties determined were cooking time and water uptake of the grain, water and oil absorption, foam capacity and stability, and emulsion capacity and stability. *Padi Tuya* was longer, broader and thicker than *Wang Kae*. The results showed both varieties to be a useful source of macronutrients. The moisture, fat and fibre in *Wang Kae* were significantly higher ($p < 0.05$) than *Padi Tuya*. However, the cowpea varieties had comparable protein levels with a mean of 17.7 g/100 g. The cooking time, hydration capacities and indices of the two varieties were also significantly different ($p < 0.05$), but their swelling capacities, indices and water uptake indices were comparable. Cooking times were 40 and 62 min for *Padi Tuya* and *Wang Kae*, respectively. The flours of the two varieties had comparable water and oil absorption capacities as well as for the swelling indices, but their emulsion capacities and stabilities were significantly different ($p < 0.05$). *Padi Tuya* had a higher foam capacity with lower stability compared to *Wang Kae*. The physico-mechanical properties determined would aid in the design of cleaning, handling and other processing machinery as well as storage facilities for the new cowpeas. The functional and the compositional parameters would assist in the food formulations.

Key words: Cowpea (*Vigna unguiculata*) varieties; compositional; physico-mechanical and functional properties



INTRODUCTION

Cowpea (*Vigna unguiculata*) is an important source of plant protein and, therefore, used in several tropical and subtropical countries all over the world. In Ghana, cowpea is mainly prepared and eaten as a whole or part of a meal. It is used in foods such as *Koose* (cowpea fritters), *gari* and beans (roasted grated fermented cassava and cooked beans) and *Tubani* (steamed bean cake). It is also used in stews and soups. These qualities make it an ideal crop for catering for the food security needs of several communities [1]. The savannah, coastal and transition zones are the major areas where cowpea is widely grown in Ghana [2]. The rainfall pattern in Ghana is such that the northern sector is characterized by little rainfall while the southern sector receives major rains. The wet conditions at the southern sector makes it non ideal for cowpea cultivation due to diseases and pest infestation. Majority of cowpea is cultivated at the northern part of Ghana. Over there, the rains fall between May and October with an average annual rainfall between 900 and 1100mm. The cowpea season is from April to July and from July to October [2, 3]. Thus, cowpeas are therefore well adapted to harsh farming systems.

Unfortunately, the current challenges with climate change in most farming systems in the tropics have brought on several biotic and abiotic factors that have affected the production, storage and even the processing of cowpea. In response to these effects, the Council for Scientific and Industrial Research (CSIR)-Savanna Agriculture Research Institute developed climate-resilient and high-yielding cowpea varieties such as, *Padi Tuya* and *Wang Kae*. *Padi-Tuya* was developed to address the challenges with low yields and long cooking times of some earlier varieties. *Wang Kae*, on the other hand, developed to be early maturing, resistant to aphids and striga infestation as well as address the issue of low yields and long cooking times.

Several researchers, Adu-Dapaah and Addison [4], David and Acka [5] and Appiah *et al.* [1] have variously worked on previous varieties of cowpea grown in Ghana, but none on these two. These two varieties could potentially be rich in nutrients, with good functionality for food application. Given the current adverse effects of climate change on the farming systems in northern Ghana, these two newly improved cowpea varieties will now be increasingly cultivated and therefore certainly play major roles in the food security systems in Ghana and for most families in West Africa. It is important that their full food-uses potential as well as key parameters needed for processing and handling machinery for these cowpeas are determined. The study therefore determined the compositional, physico-mechanical and functional properties of these newly developed cowpea varieties.

MATERIALS AND METHODS

Source of Cowpeas:

The two cowpea varieties used in this study, *Padi Tuya* and *Wang Kae*, were obtained from experimental plots of the Cowpea Improvement Program, CSIR- Savanna Agricultural Research Institute. The grains were cleaned and sorted when received before being packaged into sacks and kept at Physicochemical Laboratory of the Food



and Drugs Authority, Accra, at temperatures 28 ± 2 °C and relative humidity $75 \pm 5\%$ until ready for use.

Analyses of Physico-mechanical Properties:

Weight and volume

Both the hundred seed weight and volume were determined according to methods as explained by Wani *et al.* [7].

Bulk and true density

Bulk density (*Db*) was determined by placing 100 randomly selected seeds into a measuring cylinder and recording the volume after gently tapping the cylinder on the table for a few minutes. The ratio of mass of seed to the volume recorded gave the bulk density [7]. True density (*Dt*), which is the ratio of mass of cowpea seed to the true volume of seeds, was determined by the toluene displacement method [7]. The volume of toluene displaced was determined by immersing a weighed quantity of cowpea seed in the toluene.

Dimensions

Length (L), breadth (B) and thickness (T), were determined using a digital vernier caliper with a least count of 0.01mm (Mitutoyo, Japan). The length of the grains was determined by measuring the longest dimension (along the split of the cotyledon), while breadth and thickness were measured at the lateral and dorsal sides respectively as described by Appiah *et al.* [1].

Sphericity and surface area

The sphericity of the cowpea seeds was calculated as a function of the three principal dimensions, as explained by Wani *et al.* [7], using the formula:

$$Sphericity = \left[\frac{\sqrt[3]{LBT}}{L} \right] \times 100; \dots\dots\dots (1)$$

Where:

- L= Length of cowpea grain;
- B= Breadth of cowpea grain;
- T= Thickness of cowpea grain

The surface area was calculated using the formula:

$$Surface\ area = \frac{\pi BL^2}{2L-B} \dots\dots\dots (2)$$

Where:

- B= Breadth of cowpea grain
- L= Length of cowpea grain

2.2.5 Porosity



The Porosity of the Seeds was calculated using the formula:

$$Porosity = \left[1 - \left(\frac{Db}{Dt} \right) \right] \times 100 \dots\dots\dots(3)$$

Where:

Db = Bulk density

Dt = True Density

Angle of repose

Cowpea flour (250 μ m) was poured gradually into a funnel suspended 3 cm above a bench to form a heap on a paper at the base of the funnel. The diameter of a circle drawn around the heap was measured at four different points. Angle of repose was calculated using the formula [10]:

$$Angle\ of\ Repose = \tan^{-1} \frac{h}{0.5xd} \dots\dots\dots(4)$$

Where:

h = Height from the tip of the funnel to the bench

d = Diameter of the circle drawn

Analyses of Compositional Properties:

Compositional analyses carried out on the cowpeas were moisture, ash, crude fat and crude fibre, protein and carbohydrates. Moisture was determined according to AOAC methods 934.01 [11]. Ash by AOAC method 942.05 [11] and crude fat and fibre by AOAC [11] methods, 920.39 and 978.10, respectively. Protein was determined by the Kjeldahl method (AOAC method 988.05) where the nitrogen content determined was multiplied by the factor 6.25. Carbohydrate was estimated by difference.

Analyses of Functional properties:

The functional properties determined were partly on the grains and partly on the flour. For the grains, the parameters determined were the hydration capacity and index, swelling capacity and index, water uptake ratio and cooking time. For flour, the parameters determined were water and oil absorption, foam capacity and stability, emulsion capacity and stability, swelling power and index.

Hydration capacity and hydration index

Five gram (5 g) of cowpea was soaked in 50 mL distilled water in a measuring cylinder [12]. The seeds were left to soak for 24 h at room temperature, drained and excess water blotted with tissue paper. Thereafter, the weight of swollen seeds was measured and the hydration capacity and index were respectively calculated using the formulae:

$$\frac{Hydration\ capacity = \frac{Weight\ after\ soaking - Weight\ before\ soaking}{Number\ of\ grains} \dots\dots\dots(6)}$$



$$\text{Hydration index} = \frac{\text{Hyration capacity}}{\text{Weight of one seed}} \dots\dots\dots(7)$$

Water uptake ratio

In this method, 20 g of cowpea seeds were cooked in 200 ml of distilled water for minimum cooking time [13]. Cooked seeds were removed, drained and excess water blotted with tissue paper. The seeds were weighed and the water uptake ratio determined as a ratio of weight gained after cooking to weight before cooking.

Cooking time

Cooking time was determined according to Wani *et al.* [8] with slight modifications. In brief, 200 mL distilled water was boiled in a flask fitted with reflux condenser to prevent water loss during cooking. Thirty gram (30 g) of cowpea seeds were added to the flask and boiled for 30 min. After this period three to five seeds were withdrawn using tweezers at 2 min intervals and tested for softness by pressing between fingers and the thumb. The time from addition of seeds till attainment of a desired softness was recorded as the cooking time.

Water and oil absorption properties

Water and oil absorption capacities, foam capacity and stability were determined according to methods described by Appiah *et al.* [1], emulsion capacity and stability by Sridaran *et al.* [14], and then finally swelling power and index of the cowpea flour by methods of Appiah *et al.* [1] and Adebowale *et al.* [11], respectively.

Statistical Analysis

Quantitative variables were summarized using means and standard deviations. Student's *t*-test was carried out using Minitab Release 17 statistical software (Minitab Inc., USA) to determine differences in means at 95% confidence level.

RESULTS AND DISCUSSION

Physico-mechanical Properties

The dimensional analysis of *Padi Tuya* and *Wang Kae* showed that *Padi Tuya* is significantly ($p < 0.05$) longer, broader and thicker than *Wang Kae*, as indicated by the breadth and thickness (Figure 1).




Figure 1: Spatial dimensions of *Padi Tuya* and *Wang Kae* cowpea varieties (means with standard deviation (SD) error bars)

Again, the dimensional sizes of all these new varieties are significantly higher than those of previous Ghanaian cowpea varieties, *Nhyria*, *Tona* and *Adom*, as reported by Appiah *et al.* [1]. The lengths, breadths and thicknesses of these previous varieties are 7.6, 5.9 and 4.5 mm, respectively, for *Nhyria* variety; 7.7 mm, 5.8 mm and 4.6 mm, respectively, for *Tona* variety; and 7.7 mm, 6.3 mm and 4.9 mm, respectively, for *Adom* variety. These differences might influence the size of the screens used in rotary vibratory screens typically used for cleaning cowpeas during industrial processing [15]. Cleaning of cowpea is one of the critical unit operations during the processing of end products from cowpea. This is because cowpeas have high tendency to be contaminated with other materials such as earth, small pebbles, plant and insect waste as well as immature seeds.

An additional implication of the relatively bigger size of *Padi Tuya* could be due to the use of hermetic systems to store these cowpeas. Cowpea is usually heavily attacked by storage pests [18] and one way of controlling such pests is the use of hermetic storage, which creates a low oxygen, high carbon dioxide atmosphere which is lethal to storage insects. In hermetic storage systems, a modified atmosphere condition is created, in which the oxygen concentration is gradually reduced whilst the carbon dioxide concentration increases with time, through respiration of both the commodity and insects [20,21]. Cowpea grain size influences the movement of oxygen in hermetic containers used for cowpea grain storage. When grains are bulkier and bigger, they do not pack well in silos and creating big spaces between grains facilitating easier movement of oxygen between the bigger grains than between the smaller grains. The initial volume of oxygen in the spatial spaces will be critical to onset of the actual modified atmosphere conditions. The bigger the volume, the more time required to reach modified atmosphere. This means the time required to reach low oxygen levels in hermetic storage systems for *Padi Tuya*, through respiration effects, may be longer than in those of *Wang Kae* and earlier cowpea varieties, with relatively smaller sizes.

The 100 grain seed weight of *Padi Tuya* and *Wang Kae* were 20.54 and 18.69 g, respectively (Table 1). These were heavier compared to those of *Nhyira*, *Tona* and *Adom* varieties that were 13.17 g, 14.18 g and 15.16 g, respectively [1], and 13.08 g and 9.88 g, respectively for red and black cowpea [17]. The grain bulk and true density, grain weight and volume, are important factors to consider when designing and using industrial rotary cleaning machines for these cowpea grains [15]. The 100 grain seed weight and volume were significantly different ($p < 0.05$) between *Padi Tuya* and *Wang Kae*. Angle of repose was higher ($p < 0.05$) in *Wang Kae* than *Padi Tuya*. Previous values reported for other legumes were lower, and ranged between 17.7-19.0 ° for black gram [8], 26.6 – 31.1 for green gram [19] and 18 – 31 for pigeon pea [20]. A significantly higher porosity was observed for *Padi Tuya*, implying that this variety would hydrate easier compared to *Wang Kae* [23].

Compositional Properties

Compositional properties of the two cowpea varieties are summarized in Table 2. There was no significant difference ($p > 0.05$) between the ash, protein and carbohydrate content of the two varieties. However, fat, fibre and moisture levels of the two varieties were markedly different. The moisture content of *Padi Tuya* and *Wang Kae* in this study were comparable to *Nhyira*, *Tona* and *Adom*, which had 9.79, 9.15 and 9.83 %, respectively [1]. Otitaju *et al.* [22] reported the moisture content of *Potiskum*, *Ife-brown*, *Orarudi* and *Aloka* varieties to be 13.48, 14.5, 11.5 and 12.91%, respectively. Affrifah [23] has observed that high moisture content ($> 13\%$) before storage could lead to the development of hard-to-cook defect in cowpea. Since their moisture content were lower than 13%, the probability of these varieties developing the hard-to-cook defect is low, with *Padi Tuya* being lower than *Wang Kae* in moisture content.

The protein content of *Padi Tuya* and *Wang Kae* was comparable. These had lower protein contents compared to some varieties previously reported by Kaptso *et al.* [23], Appiah *et al.* [1], Tresina and Mohan [24] and Kaptso *et al.* [25]. The amount of protein in cowpea is important because proteins are the main molecules that imbibe water in seeds [4]. The high protein content of cowpea, has been known to reduce protein energy disorders such as *Kwashiorkor* [1]. The quality and quantity of protein in cowpea makes it a good complementary food with cereals.

Total ash for the two varieties was 3.7% and this was higher than the ash content reported for some other varieties. For instance, Appiah *et al.* [1] recorded 2.9, 3.0 and 3.2%, respectively, for *Nhyira*, *Tona* and *Adom*. The amounts of fat in *Padi Tuya* and *Wang Kae* were lower than the values reported by Appiah *et al.* [1] for *Nhyira* (2.5%), *Tona* (2.5%) and *Adom* (4.0%) varieties of cowpea. Fat content of cowpeas enhances the palatability of cowpea paste-based foods such as *Koose* or *Akara* [1, 26]. In this study, the mean fibre content of 4.6% was comparable to values reported for *Nhyira*, *Tona* and *Adom* varieties of cowpea [1]. Dietary fibre has been shown to aid digestion, regulate blood sugar level, and may play a beneficial role in diabetic patients [27].



Cooking Characteristics

Table 3 shows that *Padi Tuya* had a significantly ($p < 0.05$) shorter cooking time of 42 min compared to *Wang Kae* which required 62 min to cook to desirable texture. This indicates that *Padi Tuya* would require less energy for processing. The difference in cooking time may be attributed to differences in their hydration properties, and this may be linked to differences in their protein content [28], even though this was marginal. Additionally, the higher porosity of *Padi Tuya* seeds may have played an influential role. High hydration and swelling capacities are desirable in cowpea processing as they reduce cooking time by increasing energy transfer to the grain [28]. Cooking time of the cowpea in this study was longer than the range (24 – 42.5 min) reported by Addo *et al.* [29], comparable to some varieties (35 – 57 min) studied by Olapade *et al.* [30] and *Nhyira* (57 min) and *Tona* (65 min) reported by Appiah *et al.* [1]. Other varieties reported by Olapade *et al.* [30] and Appiah *et al.* [1] had longer cooking times of 80 min and 84 min, respectively, compared to *Padi Tuya* and *Wang Kae*. Compositional, structural and genetic variations may be responsible for those reported differences. Even though cooking improves the digestibility of proteins and carbohydrates, prolonged cooking is undesirable and may affect protein quality. Long cooking time may also lead to loss of nutrient and undesirable quality [23]. Seed swelling capacity, swelling index and water uptake ratio did not necessarily affect their cooking time. Swelling capacity and index for these two varieties were higher than 0.05 mL/seed and 0.001 reported correspondingly for these indices by Tresina and Mohan [24]. Swelling capacity was, however, lower than 2.3 – 3.2 mL/seed reported by Olapade *et al.* [30].

Functional Properties of cowpea flour

Cowpea flour is a versatile food ingredient with wide application in the processing of *Akara*, *Moinmoin* and bakery products [31]. Its performance in these products depends on the flour functionality, and this is largely dictated by protein and carbohydrate composition. In this study, results of some key functional properties evaluated are summarized in Table 4.

A mean of 1.4 and 1.1 g water/g flour was respectively recorded for water absorption capacity and oil absorption capacity in the two cowpea varieties. For these parameters, T-test showed no marked differences ($p > 0.05$) between the two cowpea varieties. Higher water absorption capacities of 1.6 – 1.9 g/g flour and 2.1 – 2.8 g/g flour have been respectively reported by Chinma *et al.* [13] and Olaofe *et al.* [32] for some Nigerian varieties. Ghanaian varieties studied by Appiah *et al.* [1] also had higher water absorption capacity (1.9 – 2.2 g/g flour) and oil absorption capacity (2.0 – 2.3 g/g flour) than the varieties used in the present study. While swelling power and true density differed significantly, the bulk densities of the cowpea were quite similar. The swelling power was higher than the 2.7 reported in a similar study [1]. The emulsion capacity of *Padi Tuya* and *Wang Kae* was approximately the same, but the latter formed a more stable emulsion. Proteins in flour aid in the formation of emulsions and, therefore, the low emulsion stability of *Padi Tuya*, could be due to the slow diffusion of the proteins to the interfacial area to stabilize the emulsion [33]. The emulsion capacity observed in this study was much lower than that observed by Hamid *et al.* [12] for red and black cowpea varieties which was 35% and 34%, respectively. The emulsion stability of the

cowpea varieties by Hamid *et al.* [12] was also higher (96% and 74%) than what was observed in this study, which could be due to differences in protein structure and functionality.

Foaming properties were significantly different ($p < 0.05$) for the two varieties, with *Padi Tuya* having higher foam capacity with less stability compared to *Wang Kae* (Figure 2).




Figure 2: Foam stability of flour of *Padi Tuya* and *Wang Kae* cowpea varieties

Foaming in cowpea is associated with the high water solubility of cowpea protein and its ability to form stable layers around gas bubbles in a liquid phase [34]. As indicated by Kethireddipalli *et al.* [35], foaming is important for the development of desirable texture in *Akara*, a product made from cowpea flour. Foaming capacity of the varieties used in this study was higher than the reported range of 10-21% [1] but within 40 – 80% for cowpea flour [33]. Variations in the functional properties of two varieties and others reported in related literature may be ascribed to compositional differences, physical properties, genetic variations and growing conditions.

CONCLUSION

The two cowpea varieties had differences in their physico-mechanical, compositional and cooking properties as well as marked differences in the functional properties of their flours. Their compositional properties indicate that they are a valuable source of macronutrients, making them potentially useful in the formulation of infant meals. While their functional properties exhibit the suitability of the two cowpea varieties in processing, the physico-mechanical properties of *Padi Tuya* and *Wang Kae* provide data that would influence grain design and operations industrial cleaning machinery.

ACKNOWLEDGEMENTS

We wish to acknowledge the assistance of technical staff at the Food Physicochemical Unit of Food and Drugs Authority of Ghana.

Declaration of Competing Interest

The authors declare that they have no interest to declare for this publication.



Table 1: Physico-mechanical properties of *Padi Tuya* and *Wang Kae* cowpea varieties

Physico-mechanical property	Cowpea varieties	
	<i>Padi Tuya</i>	<i>Wang Kae</i>
Grain Bulk density (g/cm ³)	0.72 ± 0.01 ^a	0.75 ± 0.01 ^a
Grain True density (g/cm ³)	1.13 ± 0.05 ^a	1.15 ± 0.05 ^a
Porosity (%)	35.59 ± 2.83 ^a	33.72 ± 4.26 ^a
100 Grain seed volume (cm ³)	16.67 ± 0.58 ^a	14.67 ± 0.58 ^b
100 Grain seed weight (g)	20.54 ± 0.18 ^a	18.69 ± 0.14 ^b
Sphericity (%)	72.09 ± 2.91 ^a	74.71 ± 3.43 ^b
Surface area (mm ²)	152.95 ± 15.78 ^a	145.29 ± 12.24 ^b
Angle of Repose (°)	40.24 ± 1.67 ^a	45.30 ± 1.22 ^b

*Within a row, means bearing different superscripts are significantly different (p<0.05)

Table 2: Proximate composition of *Padi Tuya* and *Wang Kae* cowpea varieties

Proximate	Cowpea varieties	
	<i>Padi Tuya</i>	<i>Wang Kae</i>
Moisture (g/100g)	10.11 ± 0.06 ^a	10.79 ± 0.14 ^b
Ash (g/100g)	3.66 ± 0.04 ^a	3.68 ± 0.08 ^a
Fat (g/100)	0.89 ± 0.03 ^a	1.51 ± 0.14 ^b
Protein (g/100g)	18.31 ± 0.77 ^a	17.16 ± 0.93 ^a
Carbohydrate (g/100g)	62.01 ± 0.08 ^a	63.17 ± 0.73 ^a
Fibre (g/100g)	5.03 ± 0.05 ^a	4.12 ± 0.23 ^b

*Within a row, means bearing different superscripts are significantly different (p<0.05)

Table 3: Cooking characteristics of *Padi Tuya* and *Wang Kae* cowpea varieties

Cooking characteristics	Cowpea varieties	
	<i>Padi Tuya</i>	<i>Wang Kae</i>
Hydration capacity (mL/seed)	0.24 ± 0.01 ^a	0.19 ± 0.01 ^b
Hydration index	0.53 ± 0.01 ^a	0.52 ± 0.01 ^b
Swelling capacity (mL/seed)	0.24 ± 0.01 ^a	0.29 ± 0.01 ^a
Swelling index	0.56 ± 0.01 ^a	0.62 ± 0.01 ^a
Water Uptake ratio	2.21 ± 0.02 ^a	2.26 ± 0.03 ^a
Cooking time (min)	42.33 ± 2.52 ^a	62.33 ± 2.52 ^b

*Within a row, means bearing different superscripts are significantly different (p<0.05)

Table 4: Functional properties of flour from *Padi Tuya* and *Wang Kae* cowpea varieties

Functional Property	Cowpea varieties	
	<i>Padi Tuya</i>	<i>Wang Kae</i>
Water absorption capacity (g/g flour)	1.42 ± 0.11 ^a	1.31 ± 0.04 ^a
Oil absorption capacity (g/g flour)	1.14 ± 0.05 ^a	1.15 ± 0.03 ^a
Swelling power	8.35 ± 0.10 ^a	7.90 ± 0.03 ^b
True density (g/cm ³)	1.39 ± 0.01 ^a	1.43 ± 0.02 ^b
Bulk density (g/cm ³)	0.87 ± 0.01 ^a	0.89 ± 0.01 ^a
Emulsion capacity (%)	6.11 ± 0.10 ^a	6.11 ± 0.96 ^a
Emulsion stability (%)	34.03 ± 3.18 ^a	47.92 ± 3.61 ^b
Foam capacity (mL)	40.00 ± 2.00 ^a	29.33 ± 1.15 ^b

*Within a row, means bearing different superscripts are significantly different (p<0.05)

REFERENCES

1. **Appiah F, Asibuo JY and P Kumah** Physicochemical and functional properties of bean flours of three cowpea (*Vigna unguiculata* L. Walp) varieties in Ghana. *Afric. J. Food Sci.* 2011; **5(2)**: 100 – 104.
2. **Akpalu MM, Salaam M, Oppong-Sekyere D and SE Akpalu** Farmers' Knowledge and cultivation of cowpea (*Vigna unguiculata* (L.) Verdc.) in three communities of Bolgatanga municipality, Upper East Region, Ghana. *Br. J. Appl. Sci. Technol.*, 2014; **4(5)**: 775 - 792.
3. **Singh BB, Asante SK, Ajeigbe H and SG Mohammed** General guide for cowpea cultivation and seed production. *Bri. J. Appl. Sci. Technol.* 2014; **4(5)**: 775 - 792.
4. **Adu-Dapaah KAH and P Addison** Variation in contents of crude protein and mineral elements in 32 cowpea (*Vigna unguiculata* (L.) Walp) accessions in Ghana. *Ghana. J. of Agric. Sci.* 2006; **39**: 83 – 86.
5. **David O and GK Acka** Comparative studies on proximate composition of cowpea, maize and soft wheat flours in Ghana. *Nutr Food Toxicol.* 2018; **3(3)**: 680 – 685.
6. **AACC 2000.** American Association of Cereal Chemists Approved Methods of the AACC, 11th ed. Method 14-22.01, Method 44-15A, St. Paul, MN: The Association.
7. **Wani IA, Sogi DS, Wani AA and BS Gill** Physical and cooking characteristics of some Indian kidney bean (*Phaseolus vulgaris* L.) cultivars. *J. Saudi Soc. of Agric. Sci.*, 2015; 1 - 9.
8. **Wani IA, Sogi DS and BS Gill** Physical and cooking characteristics of black gram (*Phaseolus mungoo* L.) cultivars grown in India. *Int. J. Food Sci. Technol.*, 2013; **48**: 255 – 2563.
9. **Tadesse FT** Handbook of farm, dairy and food machinery engineering. 3rd edition, Amsterdam, Netherlands: Elsevier Inc, 2019.
10. **Association of Official Analytical Chemists.** Official methods of analysis of AOAC International. 18th edition. Maryland, USA: AOAC International, 2006.
11. **Adebowale YA, Adeyemi IA and AA Oshodi** Functional and Physicochemical properties of flours of six mucuna species. *Afric J. Biotechnol*, 2005; **4 (12)**: 1461 - 1468.
12. **Hamid S, Muzzafar S, Wani AI and FA Masoodi** Physicochemical and functional properties of two cowpea cultivars grown in temperate Indian climate. *Cogent Food Agric.* 2015; **1**: 1 - 11.



13. **Chinma CE, Alemede IC and IG Emelife** Physicochemical and Functional Properties of Some Nigerian Cowpea Varieties. *Pak. J. Nutr.* 2008; **7(1)**: 186 - 190.
14. **Sridaran A, Karim AA and R Bhat** *Pithecellobium jiringa* legume flour for potential food applications: Studies on their physico-chemical and functional properties. *Food Chem.* 2012; **130**: 528 - 535.
15. **Irtwange SK** Design, fabrication and performance of motorized cowpea thresher for Nigerian Small-scale farmers. *African. J. of Agric. Res.*, 2009; **4 (12)**: 1383-1391.
16. **Langyintuo AS, Lowenberg-DeBoer J, Faye M, Lambert D, Ibro G, Moussa B, Kergna A, Kushwaha S, Musa S and G Ntoukam** Cowpea supply and demand in West and Central Africa. *Field Crops Res.* 2003; **82**: 215 - 231.
17. **Hamid S, Muzzafar S, Wani AI, Masoodi FA and MM Bhat** Physical and cooking characteristics of two cowpea cultivars grown in temperate Indian climate. *J Saudi Soc. Agric.* 2016; **15**: 127 - 134.
18. **Yeole NR and KC Swain** Hermetic storage technology for smallholder farmers in India. *Aayushi Int. Interdis. Res. J.*, 2018; **4 (25)**: 91 - 95.
19. **Nimkar MP and KP Chattopadhyay** Some physical properties of green gram. *J. Agric. Eng. Res.* 2001; **80**: 183–189.
20. **Baryeh EA and BK Mangope** Some physical properties of QP-38 variety pigeon pea. *J. Food Eng.* 2002; **56**: 59–65.
21. **Marabi A and IS Saguy** Effect of porosity on rehydration of dry food particulates. *J. Sci. Food Agric.* 2004; **84**: 1105–1110.
22. **Otitoju GTO, Otitiju O, Nwamarah JU and SO Baiyeri** Comparative study of the nutrient composition of four varieties of cowpea (*Vigna unguiculata*) and their products (Beans-based products). *Pak. J. Nutr.* 2015; **14 (9)**: 540 - 546.
23. **Affrifah NS** *Improving the textural properties and storage stability of cowpeas: process development and evaluation*. PhD Thesis submitted to the University of Georgia, Athens, Georgia. 2004.
24. **Kaptso KG, Njintang YN, Komnek AE, Hounhouigan J, Scher J and CMF Mbofung** Physical properties and rehydration kinetics of two varieties of cowpea (*Vigna unguiculata*) and Bambara groundnuts (*Voandzeia subterranean*) seeds. *J. Food Eng.*, 2008; **86**:91-99.
25. **Tresina PS and VR Mohan** Physicochemical and antinutritional attributes of gamma irradiated *Vigna unguiculata* (L) Walp. Subsp. *Unguiculata* seeds. *Int. Food Res. J.*, 2012; **19**:639-646.

26. **Kaptso GK, Njintang NY, Nguemtchouin MGM, Amungwa AF, Scher J, Hounhouigan J and CMF Mbofung** Characterization of morphology and structural and thermal properties of legume flours: cowpea (*Vigna unguiculata* L. Walp) and Bambara Groundnut (*Vigna subterranean* L. Verdc.) varieties. *Int. J. Food Eng.* 2015; **12**:2. <https://doi.org/10.1515/ijfe-2014-0146>
27. **Tresina PS and VR Mohan** Physicochemical and antinutritional attributes of gamma irradiated *Vigna unguiculata* (L) Walp. Subsp. *Unguiculata* seeds. *Int. Food Res. J.*, 2012; **19**:639-646.
28. **Mogos T, Dondo C and AEA Iacobini** Review of dietary fibre in the diabetic diet. *Rom. J. Diabetes Nutr. Metabol. Dis.* 2017; **24** (2): 161 - 164.
29. **Mwangwela AM** Physicochemical characteristics of conditioned and micronized cowpeas and functional properties of resultant flours. PhD Thesis submitted to University of Pretoria, South Africa 2006.
30. **Addo JK, Amoah S, Boateng AM, Okyere H, Adu-Dapaah H and A Amoah-Owusu** Physical, functional and cooking characteristics of six newly released cowpea (*Vigna unguiculata* (L) Walp) varieties in Ghana. *Agric. Food Sci. J. Ghana* 2016; **9**:742-753.
31. **Olapade AA, Okafor GI, Ozumba AU and O Olatunji** Characterization of common Nigerian cowpea (*Vigna unguiculata* L. Walp) varieties. *J. Food Eng.* 2002; **55**:101-105.
32. **McWatters KH, Ouedraogi JB, Resurreccion AVA, Hung Y-C and RD Phillips** Physical and sensory characteristics of sugar cookies containing mixtures of wheat, fonio (*Digitaria exilis*) and cowpea (*Vigna unguiculata*) flours. *Int. J. Food Sci. Technol.* 2003; **38**: 403 -410.
33. **Olaofe O, Umar YO and GO Adediran** The effect of nematicides on the nutritive value and functional properties of cowpea seeds (*Vigna unguiculata* L. Walp.). *Food Chem.* 1993; **46**: 337-341.
34. **Khalid II and SB Elharadallou** Functional properties of cowpea (*Vigna unguiculata* L Walp) and lupin (*Lupinus termis*) flour and protein isolates. *Nutr. Food Sci.* 2013; **3**:1- 6.
35. **Rangel A, Domont GB, Pedrosa C and S Ferreira** Functional properties of purified vicilins from cowpea (*Vigna unguiculata*) and pea (*Pisum sativum*) and cowpea protein isolate. *J. Agric. Food Chem.* 2003; **51**: 5792-5797.
36. **Kethireddipalli P, Hung Y-C, McWatters KH and RD Phillips** Effect of milling method (wet and dry) on the functional properties of cowpea (*Vigna unguiculata*) pastes and end product (*akara*) quality. *J. Food Sci.* 2002; **67**: 48-52.